

The 9th European Hypoxia Symposium 2018

From Molecules to Mt. Everest – From Science to Practice

Kühroint Alm, Berchtesgaden, 14th to 17th June 2018



Congress President: T. Küpper, Prof., MD, PhD (RWTH Aachen University, Germany)

Scientific committee (in alphabetical order)

T. Küpper, Prof., MD, PhD (RWTH Aachen University, Germany),
R. Lechner, MD, PhD (German Military Hospital, Ulm/Germany),
M. Tannheimer, MD, PhD (University of Ulm, Germany),
Z. Szyguła, Prof., MD, PhD, Kraków (Poland)
H.-V. Ulmer, Prof. MD, PhD (University of Mainz, Germany)

Organising committee (in alphabetical order):

B. Hartmann (RWTH Aachen University, Germany),
H. Langhof, MD (Klinik Schönsicht, Berchtesgaden/Germany),
T. Lobensteiner (German Federal Police, Kühroint, Berchtesgaden/Germany)

In cooperation of:



Sport specific performance analysis in skimountaineering: a comparison of laboratory and field tests

Bliemsrieder B.¹, Küpper T.², Lutter C.³, Schöffl I.³,
Schöffl V.³

1. Garmisch-Partenkirchen Medical Center

2. RWTH Aachen University,

3. Sozialstiftung Bamberg; corresponding author: Bernhard.

Bliemsrieder@klinikum-gap.de

Introduction: Skimountaineering is a competition sport that has gained popularity during the last years eventually leading to being accepted as a discipline at the Youth Olympic Games in Lausanne in 2020. Along with this gain in popularity, the level of performance at international competitions rose permanently and a professional and sophisticated training approach became more and more important. Most competitions are held in altitudes between 1500 m and 3500 m and therefore a considerable amount of training takes place at various degrees of hypobaric hypoxia. After having established a sport specific spiroergometry testing protocol using regular skimountaineering equipment on a treadmill during the last years, this study compares spiroergometric results between a sport specific laboratory test (LT) on a treadmill and a field test (FT) at 3100 m altitude.

Methods: 6 athletes of the German National Team (mean \pm SD, age 24.17 ± 4.22) were tested with telemetric spiroergometry (Metamax® 3B, Cortex) twice: first with an incremental step test with skies on a treadmill and four days later on snow at 3100 m altitude after three days of intermittent hypobaric hypoxia (train high, live low concept). All athletes were exposed to repetitive intermittent hypoxia during the weeks before the test. Standard spiroergometric parameters as well as Borg scale values were recorded and individual training zones were defined according to VT2 thresholds.

Results: Mean heart rate (HR) max was 192.76 ± 7.39 for LT and 180.5 ± 8.96 for FT. Mean VO₂ peak reached 71.83 ± 11.78 for LT and 54.17 ± 8.42 for FT. HR at VT2 had a mean value of 180.83 ± 7.47 for LT and 174.17 ± 7.99 for FT, with the individual difference in HR at VT2 ranging from 1 to 12 bpm.

Conclusion: While data analysis is still in progress, preliminary results demonstrate expected altitude related changes of parameters, such as the reduction of VO₂ peak. The relatively wide range of difference in HR at VT2 strongly suggests individual field testing for professional athletes rather than a generalized correction factor applied to normoxic laboratory derived thresholds.

Does Intermittent Normobaric Hypoxic Exposure Induce Pre-Acclimatisation?

Ellwood A.

Northern General Hospital, Sheffield, UK

alison.ellwood1@gmail.com

Introduction: There are a growing number of commercial gyms offering simulated-altitude training facilities and sleep chambers to aid performance in high altitude ascents and endurance events. This review specifically look at the evidence surrounding the use of intermittent normobaric hypoxia and its effect on acclimatisation.

Methods: Review of the literature was undertaken by systematic search of electronic databases including the use of key words, named author, reference scanning and citation tracking. Electronic searches of Google Scholar, EMBASE, PubMed and Web of Science were conducted.

Results: The “hypoxic dose” generated by intermittent hypoxic exposure (IHE) falls short of that required to induce endogenous erythropoietin production and erythropoiesis, thus any benefit in acclimatisation is likely to be due to other mechanisms. Repeated exposure to lower partial pressures of oxygen results in up-regulation of transcription factors such as hypoxia inducible factor 1 α . Additionally skeletal muscle oxidative metabolism is altered by exposure to hypoxia with increased glycolytic capacity – a hypoxia-induced substrate switch.

Conclusion: There have been a number of disparate studies looking at a range of physiological end-points in the study of normobaric hypoxia. Whilst these have allowed better characterisation of the physiological responses to hypoxia, to suggest that these demonstrate evidence of pre-acclimatisation is premature. Molecular changes have been described as above but the clinical ramifications of these needs clarifying. An increasing body of research suggests normobaric and hypobaric hypoxia induce different molecular and hormonal changes and these differences may be key in successfully inducing a state of pre-acclimatisation. Future research should target several key questions if normobaric hypoxia is to be safely used for recreation, rescue and military preparedness: How persistent are the physiological effects of normobaric hypoxia? To what altitude would IHE pre-acclimatise an individual? What hypoxic dose is required and how is this best delivered?

From civd curve to frostbite prevention

Gorjanc J.

St. John of God Hospital, Spitalgasse 26, 9300 ST. VEIT/GLAN, AUSTRIA; jurij.gorjanc@mf.uni-lj.si

Study 1

Purpose: To assess whether previous freezing cold injuries (FCI) would affect digit skin temperatures and rewarming rates during a follow-up cold stress test protocol.

Materials and methods: Twenty elite alpinists participated; alpinists with previous FCI requiring digit amputations (injured, INJ: n=10 total, n=8 male) were compared with ability-matched, uninjured alpinists (control, CON: n=10, all male). Digit skin temperature was measured using infrared thermography as an index of peripheral digit perfusion after a cold stress test, which consisted of 30 minutes of immersion in 8 °C water.

Results: Although, absolutely, digits in the INJ group were colder in all phases, there were no differences between the rates of rewarming of digits in both groups (Toes: INJ, $0.5^\circ \pm 0.11$ °C/min; CON, $0.7^\circ \pm 0.3$ °C/min); p=0,16). Fingers in the INJ group were colder than in the CON group after cold phase (2nd + 5th finger; p<0,05). During recovery, fingers in the INJ group were by 20% colder than in the CON group.

Conclusions: Even after FCI that requires digit amputation, there is no evidence of different tissue rates of rewarming between the injured and uninjured fingers or toes of elite alpinists.

Study 2

Purpose: The aim of the study was to assess whether previous freezing cold injury in fingers and/or toes might predispose alpinists to greater risk of further freezing cold injury, primarily due to a greater vasoconstrictor response to cold, resulting in a lower perfusion, reflected in lower skin temperature.

Materials and methods: Ten elite alpinists (N = 10; 8 male and 2 female) with amputations after freezing cold injury participated in a cold-water (8°C) immersion stress test of the hands and feet. Digit skin temperatures of amputated digits, their noninjured counterparts, noninjured digits of the affected side and noninjured digits of the corresponding side were measured. The skin temperatures were compared during three consecutive phases of the cold stress test: prewarming, cold water immersion, and passive heating.

Results: Amputated toes cooled much faster compared to their uninjured counterparts (n = 26, p < 0.001), and attained lower skin temperatures during the cold exposure test (n=26, p<0.001). Higher cooling rate was observed in all the toes on the injured limb compared with the toes on the uninjured limb (n = 40, p < 0.001). In contrast to the toes, the fingers on the injured limb after freezing cold injury were warmer during cooling phase compared to uninjured fingers (n = 15, p < 0.001).

Conclusions: The lower digit temperatures observed in affected toe stumps during the cold stress test compared to the nonamputated toes may indicate a heightened risk of future freezing cold injury with subsequent cold exposures, as a consequence of less perfusion, reflected in the lower skin temperature. This relationship was not confirmed in the fingers.

Selected possibilities to improve altitude tolerance - or: How to stay healthy at high altitudes?

Lechner R.

*Department of Anesthesiology and Intensive Care, Bundeswehr Hospital Ulm
Corresponding author: ralech@web.de*

Activities at high altitude include mountaineering, military missions, humanitarian aid missions and work stays of expatriates. In addition, rescue operations can emerge in all those fields. Therefore, it is paramount to know strategies to cope with the challenges of mountainous environment / high altitude. The three main environmental factors of mountainous environment are difficult terrain, temperature and hypobaric hypoxia.

Possibilities to counteract the challenges of mountainous environments, especially hypobaric hypoxia, are acclimatization according to international recommendations like the UIAA or the Wilderness Medical Society. In addition, preacclimatization at natural heights or in hypobaric or normobaric chambers is an alternative and increasingly used among mountaineers. However, knowledge on deacclimatization is poor. By using the latency time of altitude sickness, the impact of altitude can be reduced if people return before symptoms of high altitude sickness develop. A high physical activity at altitude, especially anaerobic exercise, is a risk factor for altitude sickness and should therefore be reduced, for example by using alternative means of transportation, having more breaks and simply reduce work load and pace. Because of susceptibility to mountain sickness is highly individual, a personnel selection of people who are not highly susceptible to hypobaric hypoxia can contribute to build an effective team. A carbohydrate rich diet, a sufficient hydration and thorough hygienic measures mitigate the impact of altitude and the hygienic constraints often found in mountainous environment. If altitude sickness occurs despite the mentioned prevention strategies a fast diagnosis and evidence-based treatment according to international recommendations is necessary. Special equipment, special training, experienced team members and a sound leadership also contribute to staying healthy at high altitude, especially by reducing the impact of environmental co-factors like UV radiation, cold, wind and alpine dangers.

Inflight measurements of pilots oxygen saturations during high-altitude glider flights of the mountain wave project (MWP) in the Himalayan region and the French Alps

Ledderhos C., Heise R., Gammel Ch., Gens A.

German Air Force Center of Aerospace Medicine

Introduction: After two expeditions to the Andes, the OS-TIV-Mountain Wave Project Team carried out a research campaign to the Himalayas in 2013/2014. Based on a motor glider-bound multidisciplinary approach, the team amongst others realized scientific measurements of the distribution and transport of atmospheric pollutants in Nepal as well as new digital aerial imageries of exceptional resolution and highest accuracy to create a 3D terrain model of glaciers close to the Mount Everest and Aconcagua. Within the project, members of the MWP worked together with scientists of the German Aerospace Center (DLR, Institute of Optical Sensor Systems), the Aerospace Technology Faculty of the University of Applied Sciences Aachen, the Karlsruhe Institute of Technology (KIT) and the German Air Force Center of Aerospace Medicine (GAF CAM) to put their concerted efforts into different objectives.

Objectives of the overall project: Firstly, participating pilots wanted to explore the Himalayas and the Mt. Everest region in pioneer flights flying up to the height of the Mt. Everest with a motor-glider (STEMME S10VT). Secondly, the project served to enhance the knowledge in turbulence research and improvement of weather forecast models. Thirdly aerosol measurements were conducted in order to record the vertical distribution of particulate matter and their optical properties for source apportionment in Nepal. Fourthly, digital aerial imagery with a resolution, as high as never used before, were taken in this region to enhance the geographical knowledge about this region, draw conclusions regarding global warming, predict flood impacts and produce documentations of natural or cultural heritage. And last but not least, concomitant aeromedical aspects of human performance and limitations with a special focus on an early detection of hypoxic hypoxia in pilots under these uniquely extreme environmental conditions (high altitude, low temperature, turbulences) were of particular interest. Historically oxygen deprivation was one of the first environmental challenges Aerospace medicine had to deal with. As recent high performance aircraft incidents have demonstrated impressively, it is still an issue this field of medicine is engaged in. Till now, sensors to detect oxygen deficiencies in flight are not in common use. With the introduction of pulse oximetry (SpO_2), an easy-to-use method to quantify oxygen deprivation of individuals was established in clinical routine. However, and this is surprising, not much effort re-

garding this technology in the field of Aerospace Medicine was made so far. Earlier work of our group of the GAF CAM could demonstrate that several of those pulse oximetry systems, used in clinical routine, are accurate enough to be also considered as warning systems for pilots. With this in mind a reflectance pulse oximetry system was tested in the environment of high-altitude glider flying. In addition, the intended measurements were tailored to optimize the crew's oxygen management during those high-altitude flights.

Methods: During the campaign the signal failure rate of pulse oximetry sensors was determined in 28 high-altitude glider flights (up to FL 290) in the Himalaya including the Mt. Everest region and (to enhance the number of flights) in the French Alps. All 7 individuals were equipped with three Nonin, bicolor reflectance sensors and the associated WristOx 3100 system. The sensors were fitted at three different measuring points (forehead, sternum and shin). In addition to oxygen saturation, the heart rate and altitude values were also recorded during the flights.

Results: In total, 19 complete data sets with readings from all sensor locations used could be gathered. SpO_2 values recorded at different measuring points showed distinct differences: the highest values were observed at the forehead ($95.39 \pm 2.11\%$) (Mean \pm standard deviation), the values detected at the sternum were the lowest ($84.38 \pm 6.74\%$). Furthermore, the failure rate was highest at the sternum ($26.51 \pm 18.56\%$ of the measuring time). At the shin, the mean SpO_2 was $93.93 \pm 4.24\%$ with a failure rate of $11.67 \pm 19.74\%$ of the measuring time. Measurements at the forehead were most robust, showing a failure rate of $3.28 \pm 6.29\%$ of the measuring time. The mean duration of failures was 12.8 ± 4.7 s at the forehead, 15.5 ± 12.8 s at the sternum and 13.7 ± 11.0 s at the shin.

Conclusions: Without any question, the use of pulse oximetry added to enhance flight safety in this campaign to the Himalayas. However, the results also clearly reveal the difficulties of obtaining robust signals that reliably reflect inflight blood oxygen saturation that have to be overcome. This is particularly true for the shortfalls anyway inherent to the method itself. Nevertheless, in this day and age, the objective continuous detection of a pilot's oxygen deprivation is inevitable with respect to flight safety. Among the application points for the sensors tested, the forehead appears to be most suitable.

Effect of endurance training conducted under hypoxia conditions, on change of exercise tolerance of patients after myocardial heart infarction (STEMI) treated with angioplasty or coronary artery bypass grafting

Nowak A.¹, Nowak Z.¹, Kucio C.¹, Küpper T.²

1. *The Jerzy Kukuczka's Academy of Physical Education, Faculty of Physiotherapy. Katowice, Poland,*

2. *RWTH Aachen University, Germany*

Corresponding author: Agata Nowak a.nowak88@gmail.com

Aim: The main aim of the study was to assess the effect of endurance training conducted under hypoxia conditions, on change of exercise tolerance of patients after myocardial heart infarction (STEMI) treated with angioplasty or coronary artery bypass grafting.

Methods: 30 patients after myocardial infarction were randomly divided into two groups: A) 12 men ($60 \pm 10,60$ years) participated in 22 training units under artificial hypoxia conditions (2000 m asl), B) Control group- 18 men ($57,16 \pm 9,72$ years) participated in 22 training units under normoxia conditions. At the beginning (in normoxia conditions) all patients underwent: spiroergometric test using a treadmill, laboratory tests: lactic acid concentration (before and after test), morphology, lipids, mass and body composition. The oxygen saturation was measured each time during every training. After 22 days of training, all groups have been re-examined (normoxia conditions)

Results: First group: after the training there were statistically significant changes in: the test time (0.001) distance (0.002), HR_{peak} (0.012), RR_{peak} - systolic (0.003), diastolic (0.030), HCT (0.033). In the case of other analyzed parameters, favorable changes were obtained, however they are not statistically significant. Second group: after the training there were statistically significant changes in: the test time (0.007), distance (0.008), MET (0.000), LA_{rest} (0.017), LA_{peak} (0.020), HR_{rest} (0.001), HR_{peak} (0.000), RR_{rest} -systolic (0.036), RR_{peak} -systolic (0.016), RBC (0.020), HCT (0.003). Comparing the two groups, differences were obtained in: MET (0.045), VE (0.002), RBC (0.012), HCT (0.006).

Conclusion: Exercise tolerance was improved in both groups. However, due to the lower oxygen content under hypoxia conditions, the range of lesions was smaller than in control group.

Pre-acclimatization in normobaric hypoxia before climbing Manaslu (8163 m)

Tannheimer M.

*Department of Sport and Rehabilitation Medicine,
University of Ulm, Leimgrubenweg 14, 89075 Ulm, Germany;
markus.tannheimer@arcor.de*

Introduction: Above 2500 m acclimatization is necessary and this requires time. The use of normobaric hypoxia allows pre-acclimatization at home instead of spending extra days on site in order to acclimatize following well established protocols [1; 2]. As a matter of fact pre-acclimatization is requested in the wide range of tourism activity and for military missions at high altitude [4-6]. Up to now no general accepted pre-acclimatization protocol is established [3].

The aim of this pilot study was to proof if pre-acclimatization in normobaric hypoxia allows a rapid ascent to high and extreme altitude and if this is practicable during normal life.

Methods: Seven healthy soldiers of the German Special Forces pre-acclimatized over 10 days using rented Hypoxic Everest Summit II generators. They adapted the hypoxic training during sleep, at rest and during physical exercise to their normal daily work routine. All persons used a more progressive acclimatization profile than that recommended by the renting company. After the 10 days of training it took 5 days including the flight to Kathmandu (1400 m) until real high altitude exposure.

Result: During normobaric hypoxia training no serious incident happened. The group was able to fly by helicopter to Samergaun (3530 m) and to walk up to the base camp (4880 m) the next day. One day later the group slept in the base camp and two more days later they climbed to camp 1 (5600 m). Nine days after leaving Kathmandu camp 3 (6850 m) was reached. None of the soldiers suffered from acute mountain sickness.

Discussion: Normobaric hypoxia enables pre-acclimatization if on-site acclimatization is not possible. During normal work routine its application during sleep is most suitable. It remains unclear if this is the most effective way for pre-acclimatization as well as the minimal time of daily hypoxic exposure needed is unknown at present. Concerning the altitude profile for pre-acclimatization our experience suggest that the recommendations of commercial companies are far too conservative.

literature available from the author

A physiological window for detecting high altitude stress

Turner R., Dal Cappello T., Malacrida S., Brugger H., Strapazzon G.

*Institute of Mountain Emergency Medicine, Eurac Research, Italy
Rachel.Turner@eurac.edu*

Introduction: Acute high altitude exposure in healthy subjects is commonly associated with complex pathophysiological acute mountain sickness (AMS) symptomatology. Specifically, failure to adequately adapt to hypobaric hypoxia during rapid ascent to high altitude can result in hypoxemia and tissue oxidative stress (OxS). However, the non-invasive mechanistic processes necessary to define causal relationships between AMS, potential cerebral edema and increased intracranial pressure (ICP) remain unclear. Plus, the validity of potential surrogate measures for accurate definition of adaptive or maladaptive response to hypoxia remains under investigation.

Methods: A prospective study to monitor human physiological parameters on acute hypobaric hypoxic exposure of 16 healthy individuals to 3830m was conducted via direct helicopter ascent to Mount Ortles glacier. Neurological signs and symptoms of AMS, HACE, clinical parameters, ultrasonography of ONSD and biological samples were prospectively assessed at baseline (262m) and after passive ascent to 3830m (9, 24 and 72h) by blinded investigators. Reactive oxygen species (ROS) production rate was determined by means of an electron paramagnetic resonance method. OxS biomarkers were assessed by immune and/or enzymatic methods.

Results: ONSD increased with exposure to altitude in all participants ($p= 0.003$), more so in those with AMS ($p < 0.001$). After 24h the imbalance between ROS production (+141%) and scavenging (-41%) reflected an increase in OxS related damages by 50%-85% ($p < 0.05$). Despite the concurrent increase of ONSD, regression analysis did not infer a causal relationship between OxS biomarkers and changes in ONSD.

Conclusion: The relationship between ONSD, time, and mode of ascent to high altitude is extremely relevant for the delineating the role of sonography in supporting a diagnosis of AMS. These results provide new insight regarding ROS homeostasis and potential pathophysiological mechanisms of acute exposure to hypobaric hypoxia. However, caution should be recommended, as there remain limitations to the diagnostic utility of ONSD as a surrogate measure of ICP in these environments. In addition, mechanistic investigations are warranted to identify the sources of ROS production, the sequence of signaling pathways involved and timing of concomitant upregulation of biomarkers of OxS. Future work should focus on large cohort

studies, where a range of concurrent environmental stressors may be investigated under controlled, standardised, hypobaric hypoxic conditions.

Sports Anemia and Hypobaric Hypoxia: Clinical and Biochemical Evaluation of Hematologic Changes and Muscle Damage during the Ultramarathon Aconcagua

Schmidt A.¹, Acuña A.¹, Borzzota F.¹, Duplessis R.¹, Fiorentino Y.¹, González Costa R.¹, González Ochoa A.¹, Marengo A.¹, Maroto C.¹, Nanon E.¹, Ponce R.¹, Rogé I.¹, Saravia A.¹, Senatra P.¹, Straniero M.¹, Der Parsehian S.², Ortiz Naretto Á.³, Pereiro M.⁴, Donato S.¹

1. Andean Association for Altitude Medicine

2. Hospital Materno Infantil Ramón Sarda

3. Hospital Dr. Francisco J. Muñiz, 4. Hospital Interzonal de Agudos Pedro Fiorito

e-mail: astridschmidt@fcm.uncu.edu.ar

Introduction: Sports anemia and muscle damage have been observed especially in athletes who perform resistance exercises. The aim of this study was to evaluate in a clinical and biochemical way the reaction to hypobaric hypoxia as an additional stress factor, observing clinical status and biochemical profile as inflammatory response, production of hemolysis or rhabdomyolysis during extreme exercise in the participants of Ultramarathon Aconcagua. The competition was held on November 29 in 2014 in Aconcagua State Park (Mendoza, Argentina) with a distance of 50km and a cumulative altitude difference of 1590 meters, starting at 2900 masl, reaching 4200 masl and back to 2900 masl.

Methods: Controls of pulse oximetry, heart rate, clinical status and Lake Louise Score at Confluence base camp (3400 masl) were performed on 14 competitors. Blood, urine and saliva samples were taken the day before and two hours after the race. A questionnaire about previous mountaineering experience, allergic response (aqua test), and fatigue/anxiety/sleep quality visual scales were also done to the competitors before and after the competition.

Results: The increased values of LDH, CPK, myoglobin, troponin T ultrasensitive, NT-proBNP and White bloodcell count (neutrophils) after the race were statistically significant, but there were no changes related to the Red cells series. Saliva samples showed neither inflammatory nor immunologic changes. No significant difference was found between men and women.

Conclusion: Performing aerobic exercise in hypobaric hy-

poxia conditions is exhausting and causes subclinical muscle damage, but it is not associated with acute anemia. We consider that this study contributes to confirm that practicing this type of extreme sports does not cause pathologic or unhealthy consequences in healthy persons.

Keywords: marathon, anemia, hemolysis, rhabdomyolysis, altitude, hypobaric hypoxia.

Physiological Changes in Participants of an Ultramarathon in Subarctic Climate

Steinach M¹, Coker RH², Maggioni AM^{1,3}, Rundfeldt LC¹, Schalt A¹, Gunga HC¹

1. Center for Space Medicine and Extreme Environments
Berlin, Institute of Physiology, Charité Berlin, Germany

2. Institute of Arctic Biology, University of Alaska-Fairbanks,
Fairbanks, Alaska, USA

3. Department of Biomedical Sciences for Health, University of
Milan, Italy

The knowledge about physiological changes in athletes participating in ultramarathons is scarce. While tens of thousands athletes compete in classic marathons, only a few hundred do so in ultramarathons, i.e. distances longer than 42.195 km. Nevertheless, the number of participants continually increased over last few decades [1] and furthermore, the number and the performances of athletes participating in extreme environments are increasing as well [2]. Therefore, and in order to gain more knowledge in this field, we conducted first studies in participants of the "Yukon Arctic Ultra" (YAU). This run takes place biennially in February in northwestern Canada and covers a maximum distance of 690 km. Temperatures range around -20°C during the day and can drop to -50°C during the night (max temp during the study: +5°C, min temp: -50°C). We surveyed n=24 athletes (17 male and 7 female, age: 41.4 ± 9.6 years, BMI: 24.0 ± 2.6 kg/m²), who competed in the foot-race-category, over three years (2013 n=6, 2015 n=8 and 2017 n=10) regarding various parameters. In total, 12 of the 24 participants (50%) completed the YAU's full distance of 690 km and 14 (58%) at least 480 km. We found considerable changes among various parameters in response to the race's stress, such as in body weight, body composition, serum parameters resembling tissue damage and heart-stress as well as in parameters indicating an increased function of brown adipose tissue [3] as well as of changes in vegetative control and psychometric measures such as mood and fatigue [4].

References

[1] Whyte G (2014) Age, sex and (the) race: gender and geriatrics in the ultra-endurance age. *Extrem Physiol Med* 3: 1.

[2] Knoth C, Knechtle B, Rust CA, Rosemann T, Lepers R (2012) Participation and performance trends in multistage ultramarathons-the 'Marathon des Sables' 2003-2012. *Extrem Physiol Med* 1: 13.

[3] Coker RH, Weaver AN, Coker MS, Murphy CJ, Gunga HC, Steinach M (2017) Metabolic Responses to the Yukon Arctic Ultra: Longest and Coldest in the World. *Med Sci Sports Exerc* 49: 357-362.

[4] Rundfeldt LC, Maggioni MA, Coker RH, Gunga HC, Riveros-Rivera A, Schalt A, Steinach M (2018) Cardiac Autonomic Modulations and Psychological Correlates in the Yukon Arctic Ultra: The Longest and the Coldest Ultramarathon. *Front Physiol* 9: 35.

Nightly hypoglycaemia, an often not sufficiently known special risk for diabetics (type 1) after intensive physical activities

Ulmer H.V., Pisarz S.

Inst. Sport Science Univ. Mainz; ulmer@uni-mainz.de

Introduction: Exercise is good for diabetics, an often used recommendation. Exercise is a complication for diabetics, however, because carbohydrate supply, insuline dosage and metabolism are to be balanced. Mountaineering/hiking is performed with high physical intensities, often used only few days/year (problem of experience) with depletion of glycogen stores, filled up during sleep with the risk of unnoticed dangerous hypoglycaemia. Informal remarks led to the assumption, that this fill-up effect, especially after activities in the afternoon or later, is not sufficiently known, in context with the dead-in-bed syndrome, too: A literature survey of Tattersall & Gill (1991, [1]) mentioned not a relationship to intensive physical activities at afternoon or later. A comment from 2014 [2] focussed to cardiac aspects, too, but mentioned „hence, some deaths may have been due to other conditions such as cardiomyopathy or diabetic precoma“. An actual guideline [3] recommends typically physical activities, but describes not the special risk of nightly unnoticed hypoglycaemia after physical activities at afternoon or later in its chapter „Konsequenzen für sportliche Aktivität bei Typ-1-Diabetes“.

Methods: To clarify these aspects, we made a literature survey and interviews with 14 diabetes advisers [4].

Results: (state 2013) Scientific literature at 31% (of n=39) and additional sources at 23% (of n=35) described this effect. Nightly check of bloodglucose was recommended in literature (23%) and only at 3% in sources. In training materials (n=9) this special aspect was described twice (22%). 69% of the 14 diabetes-advisers informed explicitly about the special risk of nightly hypoglycaemia after intensive metabolic activities and 67% recommended a nightly bloodglucose check.

Conclusions: We assume, that still today the special risk of nightily hypoglycaemia after glucose-consuming activities is not sufficiently enogh well known (except the diabetes advisers). A relationship with dead-in-bed syndrome is probable and should regarded in future. Especially in context with physical activities at afternoon or later hikers and mountaineers are endagered, if they are beginners or active only in holidays (less experience in handling bloodsuger in unusual situations). As one prophylactic mean we recommend to use an alarm mobile/clock for checking bloodglucose at about 3 a.m.

literature – please ask: ulmer@uni-mainz.de

References

- [1] Tattersall R.B., Gill G.V., (1991): Unexplained deaths of type 1 diabetic patients, *diabetic medicine*, 8(1), PP. 49-58. Abstract and German short version in:
https://www.medtronic-community.de/wp-content/uploads/3-Tattersall-und-Gill_Tod-in-Bett-Syndrom.pdf
- [2] DIAPEDIA – open access knowledge base (2014): Dead-in-bed syndrome: <https://www.diapedia.org/acute-and-chronic-complications-of-diabetes/7105157816/dead-in-bed-syndrome> (state 4.7.2018)
- [3] Esefeld K., Zimmer P., Stumvoll M., Halle M. (2017): Diabetes, Sport und Bewegung. https://www.deutsche-diabetes-gesellschaft.de/fileadmin/Redakteur/Leitlinien/Praxisempfehlungen/2017/dus_2017_S2_Praxisempfehlungen_3974253_Halle_Diabetes__Sport_und_Bewegung__13__Online-PDF.PDF
- [4] Pisarz S. **Nächtliche Hypoglykämie nach Ausdaueraktivitäten bei Diabetikern Typ 1 – ein vernachlässigtes Problem?** Diploma thesis, Mainz, submitted: 13.6.2013. Abstract in: <http://www.sport.uni-mainz.de/physio/pdf/PISTARZ-DiplKurz-4Homepage21VI2013.pdf>

Exertion programming for hiking uphill by percentage guide signs

Ulmer H.-V.¹, Ringleb F.¹, Langhof H.²

1. Inst. Sport science Univ. Mainz

2. Mittendorff-Institute Berchtesgaden;

ulmer@uni-mainz.de

Intorduction: Usually guide signs for hikers in mountain landscapes indicate distances until the goal in hours. Fit hikers need less, recreation hikers more time. Especially hikers with

reduced fitness need informations about the degree of exertion essential for reaching an uphill goal or informations for turning back at time to avoid unwelcome fatigue/exhaustion. For arthritic hikers (problems in descending) a funicular is favourable.

Methods: The ascent (+ 470 m) to the attractive viewpoint Hirschkaser (1385 m, Region Ramsau, Germany), was marked by signs with 20, 40, 60, 80% of total essential exertion for reaching the goal, tested before by a very experienced hiker. At starting point special flyers explained the meaning of these percentage-signs. 126 hikers (73 m, 53 f, 27 to 85 years) were interviewed at goal about their experience in hiking, knowledge of the track and general comments about the new guide signs.

Results: 74 % were vacationists, often with previous knowledge of the track. Only 16 % used the signs for arrangement of exertion, but for 67 % the signs were assessed as a positive method for improving the attractivity of the track. As consequence the Tourismus-Information Ramsau (1st mountaineering village in Germany) printed for 2017 1000 and for 2018 2000 flyers.

Conclusions: Caused by the random conditions of the questioning mostly tourists were interviewed not elonging to the target group of recreation hikers or those with reduced fitness. The results indicate attentiveness and attractiveness of the percentage guide signs, however. So we recommend these special markings as supporting a gentle tourismus for hosts seeking recreation even with reduced fitness.

References

- [1] Ringleb, F. (2016): Attraktivität eines anstrengungsbezogenen, prozentmarkierten Wanderwegs für ambitionierte Bergwanderer und Urlaubstouristen im Berchtesgadener Land – Eine Befragung am Aussichtspunkt Hirschkaser. Bachelor thesis Mainz, abstract: <http://www.sport.uni-mainz.de/physio/pdf/PISTARZ-DiplKurz-4Homepage21VI2013.pdf>
- [2] Ulmer, H.-V. (1996): Concept of extracellular regulation of muscular metabolic rate during heavy exercise of awake men by psychophysiological feedback. *Experientia* 52, 416 - 420

Thanks to Mittendorff-Institut, Alpines Gesundheitsforum Berchtesgaden and Dr. H. Langhof (Berchtesgaden) sowie F. Rasp (Ramsau)